



Advancement

Medical Weight Loss Solutions
for Clinics and Practice Use Only

Semaglutide (7.92MG)

(OZEMPIC / WEGOVY)

\$99

Vitamin B12 0.5mg/ml
L-Carnitine 100mg/ml
Subcutaneous injection

Dosage Guide How Much to Draw from Vial

- Weeks 1-4: 0.27 mg per dose = 0.1 ml = 10 IU ON SYRINGE.
- Weeks 5-8: 0.54 mg per dose = 0.2 ml = 20 IU ON SYRINGE.
- Weeks 9-12: 1.08 mg per dose = 0.4 ml = 40 IU ON SYRINGE.
- Weeks 13-15: 2.16 mg per dose = 0.8 ml = 80 IU ON SYRINGE.
- After week 15: 2.65 mg per dose = 1 ml = 100 IU ON SYRINGE.



Dosage Guide How Much to Draw from vial

- Weeks 1-4: 2.5 mg per dose = 0.15 ml = 15 IU ON SYRINGE.
- Weeks 5-12: 5 mg per dose = 0.33 ml = 34 IU ON SYRINGE.
- Weeks 13-20: 10 mg per dose = 0.66 ml = 66 IU ON SYRINGE.
- After week 20: 15 mg per dose = 1 ml = 100 IU ON SYRINGE.

Retatrutide (45MG)

(Mounjaro / Zepbound)

\$175

Vitamin B12 0.5mg/ml
L-Carnitine 100mg/ml
Subcutaneous injection

Retatrutide (36MG)



\$450

Vitamin B12 0.5mg/ml
L-Carnitine 100mg/ml
Subcutaneous injection

Dosage Guide How Much to Draw from vial

- Week 1-4: Start with 2.5 mg once
- weekly. Inject 0.2 ml or 20 units from the syringe. Week 5-8: Increase the dose to 5 mg once
- weekly. Inject 0.4 ml or 40 units from the syringe.
- Week 9-12: Increase the dose to 7.5 mg once weekly. Inject 0.6 ml or 60 units from the syringe
- Week 13-16: Increase the dose to 10 mg once weekly. Inject 0.8 ml or 80 units from the syringe.
- Week 17 and beyond: Increase the dose to 12 mg once weekly, which is the maintenance dose.
- Inject 1 ml or 100 units from the syringe.



DELIVERY IN COLD PACKAGING

THE PACK INCLUDES



Multidose 3mls Vial semaglutide 2.64 mg/ml OR TIRZEPATIDE 15 MG/ML OR
RETATRUTIDE 12MG/ML (3ML) PLUS Vitamin B12 0.5mg/ml, L-Carnitine 100mg/ ml.
Patient full instructions on usage.
Sub cutaneous syringes.
Alcohol Wipes.

How To Place Order

We look forward to assisting you and Providing high-quality support to meet your clinic's needs.

www.coretrustlabs.com

Email: info@coretrustlabs.com

The compounded preparations referenced in this communication are formulated and produced in compliance with Good Manufacturing Practices (GMP) standards in internationally recognized facilities. These products are intended exclusively for professional use by licensed physicians. The contents herein are for informational purposes only and do not constitute an endorsement or recommendation for any specific treatment or therapy. It is the responsibility of the prescribing physician to ensure that the use of these compounded preparations is appropriate for the individual patient and to adhere to all applicable laws and regulations governing their practice. The Manufacturer assumes no liability for misuse or misapplication of products.